arXiv:1102.1878v1 [physics.gen-ph] 9 Feb 2011

A Note on Parallel Transportation in Symmetric TeleParallel Geometry

Muzaffer Adak
Department of Physics, Faculty of Arts and Sciences,
Pamukkale University, 20017 Denizli, Turkey
`madak@pau.edu.tr`

Abstract

I discuss autoparallel curves and geodesics in the symmetric teleparallel geometry.

PACS numbers: 04.50.Kd,

Keywords: Non-Riemannian geometry,

1 Introduction

We know that the autoparallel curves in the Riemann geometry coincide with its geodesics. In this note my aim is to investigate if there is a such result in the symmteric teleparallel geometry (STPG).

2 Mathematical Preliminaries

Spacetime, in general, is denoted by $\{M, g, \nabla\}$ where M is orientable and differentiable manifold, $g = g_{\alpha\beta} e^\alpha \otimes e^\beta$ is the metric tensor written in terms of 1-forms e^α and ∇ is connection associated with connection 1-forms ${\Lambda^\alpha}_\beta$. Cartan structure equations define nonmetricity 1-forms, torsion 2-forms and curvature 2-forms, respectively

$$
\begin{aligned}
Q_{\alpha\beta} &:= -\frac{1}{2}\mathrm{D}g_{\alpha\beta} = \frac{1}{2}(-\mathrm{d}g_{\alpha\beta} + \Lambda_{\alpha\beta} + \Lambda_{\alpha\beta}) \; , &(1)\\
T^\alpha &:= \mathrm{D}e^\alpha = \mathrm{d}e^\alpha + {\Lambda^\alpha}_\beta \wedge e^\beta \; , &(2)\\
{R^\alpha}_\beta &:= \mathrm{D}{\Lambda^\alpha}_\beta := \mathrm{d}{\Lambda^\alpha}_\beta + {\Lambda^\alpha}_\gamma \wedge {\Lambda^\gamma}_\beta &(3)
\end{aligned}
$$

where d and D are exterior derivative and covariant exterior derivative. Geometrically, the nonmetricity tensor measures the deformation of length and angle standards during parallel transport. For example, after parallel transportation of a vector along a closed curve, the length of the final vector may be different from that of the initial vector. On the other hand, torsion relates to the translational group. Moreover it is sometimes said that closed parallelograms do not exist in spacetime with torsion. Finally, curvature is related to the linear group. That is, when a vector is parallel transported along a closed loop, the vector undergos a rotation due to curvature. These quantities satisfy Bianchi identities:

$$
\begin{aligned}
\mathrm{D}Q_{\alpha\beta} &= \frac{1}{2}(R_{\alpha\beta} + R_{\beta\alpha}) \; , &(4)\\
\mathrm{D}T^\alpha &= {R^\alpha}_\beta \wedge e^\beta \; , &(5)\\
\mathrm{D}{R^\alpha}_\beta &= 0 \; . &(6)
\end{aligned}
$$

Full connection 1-forms can be decomposed uniquely as follows [1],[2],[3]:

$$
{\Lambda^\alpha}_\beta = \underbrace{(g^{\alpha\gamma}\mathrm{d}g_{\gamma\beta} + {p^\alpha}_\beta)/2 + {\omega^\alpha}_\beta}_{Metric} + \underbrace{{K^\alpha}_\beta}_{Torsion} + \underbrace{{q^\alpha}_\beta + {Q^\alpha}_\beta}_{Nonmetricity} \qquad (7)
$$

where ${\omega^\alpha}_\beta$ Levi-Civita connection 1-forms

$$
{\omega^\alpha}_\beta \wedge e^\beta = -\mathrm{d}e^\alpha \; , \qquad (8)
$$

$K^\alpha{}_\beta$ contortion tensor 1-forms,

$$K^\alpha{}_\beta \wedge e^\beta = T^\alpha \ , \tag{9}$$

and anti-symmetric 1-forms

$$q_{\alpha\beta} = -(\iota_\alpha Q_{\beta\gamma})e^\gamma + (\iota_\beta Q_{\alpha\gamma})e^\gamma \ , \tag{10}$$
$$p_{\alpha\beta} = -(\iota_\alpha \mathrm{d}g_{\beta\gamma})e^\gamma + (\iota_\beta \mathrm{d}g_{\alpha\gamma})e^\gamma \ . \tag{11}$$

This decomposition is self-consistent. To see that it is enough to multiply (7) from right by e^β and to use definitions above. While moving indices vertically in front of both d and D, special attention is needed because $\mathrm{d}g_{\alpha\beta} \neq 0$ and $\mathrm{D}g_{\alpha\beta} \neq 0$. Symmetric part of the full connection comes from (1)

$$\Lambda_{(\alpha\beta)} = Q_{\alpha\beta} + \frac{1}{2}\mathrm{d}g_{\alpha\beta} \tag{12}$$

and the remainder is anti-symmetric part

$$\Lambda_{[\alpha\beta]} = \frac{1}{2}p_{\alpha\beta} + \omega_{\alpha\beta} + K_{\alpha\beta} + q_{\alpha\beta} \ . \tag{13}$$

It is always possible to choose orthonormal basis 1-forms which I denote $\{e^a\}$. Then the metric $g = \eta_{ab}e^a \otimes e^b$ where $\eta_{ab} = \mathrm{diag}(\pm 1, \cdots, 1)$. In this case the splitting of the full connection (7) takes the form

$$\Lambda^a{}_b = \omega^a{}_b + K^a{}_b + q^a{}_b + Q^a{}_b \ . \tag{14}$$

If only $Q_{\alpha\beta} = 0$, connection is metric compatible; Einstein-Cartan geometry. If both $Q_{\alpha\beta} = 0$ and $T^\alpha = 0$, connection is Levi-Civita; pseudo-Riemannian geometry. If $R^\alpha{}_\beta = 0$ and $Q_{\alpha\beta} = 0$, it is called teleparallel (Weitzenböck) geometry. If $R^\alpha{}_\beta = 0$ and $T^\alpha = 0$, it is called symmetric teleparallel geometry (STPG).

3 STPG

In STPG only nonmetricity tensor is nonzero:

$$Q_{\alpha\beta} \neq 0 \quad , \quad T^\alpha = 0 \quad , \quad R^\alpha{}_\beta = 0 \ . \tag{15}$$

Here I argue that in a well-chosen coordinate (or gauge) every metric has its own nonmetricity. This can be seen a kind of gauge fixing. Let us show this argument as follows. First one has to choose the natural frame $e^\alpha = \mathrm{d}x^\alpha$ and the connection as $\Lambda^\alpha{}_\beta = 0$, then automatically $R^\alpha{}_\beta = 0$ and $T^\alpha = 0$ and $Q_{\alpha\beta} = -\frac{1}{2}\mathrm{d}g_{\alpha\beta}$. This sequence of the operations corresponds to $\omega_{ab} + q_{ab}$ in the orthonormal frame. Besides, in this geometry identities (4)-(6) give one nontrivial identity, $\mathrm{D}Q_{\alpha\beta} = 0$. From now on let Greek indices denote natural (holonomic) ones.

3.1 Autoparallel Curves

In the Riemann geometry one requires that the tangent vector to the autoparallel curve points in the same direction as itself when parallel propagated, and demand that it maintains the same length,

$$\mathrm{D}T^\alpha = 0 \; . \tag{16}$$

On the other hand, since intuitively autoparallel curves are "those as straight as possible" I do not demand the vector to keep the same length during parallel propagation in STPG. It is known that nonmetricity is related to length and angle standards at parallel transportation. Therefore I prescribe the parallel propagation of the tangent vector

$$\mathrm{D}T^\mu = (aQ^\mu{}_\nu + bq^\mu{}_\nu)T^\nu + cQT^\mu \tag{17}$$

where $T^\mu = \frac{dx^\mu}{d\tau}$ is the tangent vector to the curve $x^\mu(\tau)$ with an affine parameter τ and a, b, c are arbitrary constants. Here since I choose $\Lambda^\alpha{}_\beta = 0$ I obtain $\mathrm{D}T^\mu = \mathrm{d}T^\mu = (\partial_\alpha T^\mu)e^\alpha$. Moreover, I write $Q^\mu{}_\nu = Q^\mu{}_{\nu\alpha}e^\alpha$ and $q^\mu{}_\nu = q^\mu{}_{\nu\alpha}e^\alpha$ and $Q = Q^\nu{}_{\nu\alpha}e^\alpha$. Then Eqn(17) turns out to be

$$\partial_\alpha T^\mu = (aQ^\mu{}_{\nu\alpha} + bq^\mu{}_{\nu\alpha})T^\nu + cQ^\nu{}_{\nu\alpha}T^\mu \; . \tag{18}$$

Now after multiplying this with T^α I write $T^\alpha\partial_\alpha := \frac{d}{d\tau}$. Thus I arrive at

$$\frac{d^2x^\mu}{d\tau^2} = (aQ^\mu{}_{\nu\alpha} + bq^\mu{}_{\nu\alpha})\frac{dx^\nu}{d\tau}\frac{dx^\alpha}{d\tau} + cQ^\nu{}_{\nu\alpha}\frac{dx^\mu}{d\tau}\frac{dx^\alpha}{d\tau} \; . \tag{19}$$

Here by using $Q_{\alpha\beta} = -\frac{1}{2}\mathrm{d}g_{\alpha\beta}$ I get $Q^\mu{}_{\nu\alpha} = -\frac{1}{2}g^{\mu\beta}(\partial_\alpha g_{\beta\nu})$ and then $q^\mu{}_{\nu\alpha} = -\frac{1}{2}g^{\mu\beta}(\partial_\nu g_{\alpha\beta} - \partial_\beta g_{\alpha\nu})$. Consequently, I express the autoparallel curve equation in terms of the metric as follows

$$\begin{aligned} \frac{d^2x^\mu}{d\tau^2} &= g^{\mu\beta}\left[-\frac{a+b}{4}(\partial_\alpha g_{\beta\nu} + \partial_\nu g_{\beta\alpha}) + \frac{b}{2}(\partial_\beta g_{\alpha\nu})\right]\frac{dx^\nu}{d\tau}\frac{dx^\alpha}{d\tau} \\ &\quad -\frac{c}{2}g^{\nu\beta}(\partial_\alpha g_{\beta\nu})\frac{dx^\mu}{d\tau}\frac{dx^\alpha}{d\tau} \end{aligned} \tag{20}$$

where I symmetrized the first term of the first line in $(\nu\alpha) = \frac{1}{2}(\nu\alpha + \alpha\nu)$ indices. If I fix $a = b = 1$ and $c = 0$, then this equation becomes the same as the autoparallel curve of the Riemann geometry.

3.2 Geodesics

Intuitively, geodesics are "curves as short as possible". Interval between infinitesimal points is given by metric

$$ds^2 = g_{\mu\nu} dx^\mu dx^\nu \tag{21}$$

Let me parameterize the curve between endpoints as $x^\mu = x^\mu(\tau)$, then I obtain

$$s = \int_{\tau_1}^{\tau_2} \left(-g_{\mu\nu}\dot{x}^\mu \dot{x}^\nu\right)^{1/2} d\tau \tag{22}$$

where dot denotes τ-derivative. I inserted a minus sign because of the Lorentz signature. I wish now to derive the condition on a curve which makes it extremize the length between its endpoints, i.e., wish to find those curves whose length does not change to first order under an arbitrary smooth deformation which keeps the endpoints fixed. This condition gives rise to the Euler-Lagrange equations

$$\frac{d}{d\tau}\left(\frac{\partial L}{\partial \dot{x}^\alpha}\right) - \frac{\partial L}{\partial x^\alpha} = 0 \tag{23}$$

of the action integral $I = \int_{\tau_1}^{\tau_2} L(x^\mu, \dot{x}^\mu, \tau)$. Thus in my case the Lagrangian is

$$L = [-g_{\mu\nu}(x)\dot{x}^\mu \dot{x}^\nu]^{1/2} \tag{24}$$

where x stands for coordinate functions x^μ. Now Euler-Lagrange equations yield

$$\ddot{x}^\beta + \frac{1}{2} g^{\alpha\beta}(\partial_\mu g_{\alpha\nu} + \partial_\nu g_{\alpha\mu} + \partial_\alpha g_{\mu\nu})\dot{x}^\mu \dot{x}^\nu = \frac{\dot{x}^\beta}{2(g_{\mu\nu}\dot{x}^\mu \dot{x}^\nu)} \frac{d(g_{\mu\nu}\dot{x}^\mu \dot{x}^\nu)}{d\tau} \tag{25}$$

Attention to the last term! Let me evaluate it. First, I write it as $g_{\mu\nu}\dot{x}^\mu \dot{x}^\nu = g_{\mu\nu}T^\mu T^\nu = T_\mu T^\mu$. Now,

$$\begin{aligned} \mathrm{d}(T_\mu T^\mu) &= (\mathrm{D}g_{\mu\nu})T^\mu T^\nu + 2g_{\mu\nu}T^\mu(\mathrm{D}T^\nu) \\ &= -2Q_{\mu\nu}T^\mu T^\nu + 2g_{\mu\nu}T^\mu(\mathrm{D}T^\nu) \end{aligned} \tag{26}$$

Here the usage of Eqn(17) gives

$$\mathrm{d}(T_\mu T^\mu) = 2(a-1)Q_{\mu\nu}T^\mu T^\nu + 2cQT_\mu T^\mu \tag{27}$$

This means that if I choose $a = 1$ and $c = 0$, Eqn(25) becomes the same as the geodesic equation of the Riemann geometry.

4 Result

Thus if in STPG the parallel propagation of the tangent vector $T^{\mu} = \frac{dx^{\mu}}{d\tau}$ to a curve $x^{\mu}(\tau)$ is defined as

$$\mathrm{D}T^{\mu} = (Q^{\mu}{}_{\nu} + q^{\mu}{}_{\nu})T^{\nu} \tag{28}$$

then the autoparallel curve equation is obtained as

$$\frac{d^2x^{\mu}}{d\tau^2} + \frac{1}{2}g^{\mu\beta}\left(\partial_{\alpha}g_{\beta\nu} + \partial_{\nu}g_{\beta\alpha} - \partial_{\beta}g_{\alpha\nu}\right)\frac{dx^{\nu}}{d\tau}\frac{dx^{\alpha}}{d\tau} = 0\,. \tag{29}$$

which is the autoparallel curve equation of the Riemann geometry. Also it is shown that this is the geodesic equation of the STPG like in the Riemann geometry.

Acknowledgement

The author thanks to F W Hehl for stimulating criticisms.

References

[1] Hehl F W *et al.*, *Phys. Rep.* **258** (1995) 1

[2] Tucker R W and Wang C, *Class. Quant. Grav.* **12** (1995) 2587

[3] Adak M, *Non-Riemannian Theories of Gravitation and the Dirac Equation*, Lambert Academic Pub., 2010, ISBN: 978-3-8383-5139-1